



Praziquantel

AFFIX PATIENT IDENTIFICATION LABEL HERE

UMRN:
Family Name:
Given Name:

Medicine name	Strength	Example of brand	Dose directions



English

Praziquantel

Why do I need to take Praziquantel?

Your blood test shows that you have an infection caused schistosomiasis or bilharzia. This is a parasite that is carried by freshwater snails and is found in many countries in Asia, South America and Africa. Most people with schistosomiasis do not have any symptoms and may have been infected many years ago. If it is not treated the infection can cause liver, kidney, or bladder problems.

Before taking this medication, you should tell your doctor or nurse practitioner if you are taking any other medications. For example, medicines used to treat epilepsy, TB and HIV can interact with this medication.

You should also tell your doctor or nurse if you have had seizures or an abnormal heart rhythm in the past.

How do I take Praziquantel?

The dose of this medication has been calculated according to your weight. You need to take two doses that should be taken four hours apart.

The medicine should be taken with food. Do not chew the medication.

Do not drink grapefruit juice whilst you are taking this medication.

Do not drive or operate dangerous machinery when taking this medication.

Karen

Praziquantel

မိုးယကဘဉ်အီ Praziquantel ဒ်လဲဉ်.

နသ့ဉ်အတၢ်မၤကွၢ်ဒုးနဲဉ်ဝဲဒဉ်လၢနအိဉ်ဒီးတၢ်ဘဉ်ကူ ဘဉ်ကလၢအကဲထီဉ်ခီဖျိ schistosomiasis မ့တမ့ၢ် bilharzia န့ဉ်လီၤ.ထိးကလဲာ်တခါအံၤထံဆါအချီဉ် တဖဉ်စိာ်ဃာ်ဒီးချီဉ်တဖဉ်အံၤတၢ်ထံဉ်အီၤလၢအ့ရှၢဉ်, အမဲရကၤဒီးအဖံဉ်ဂြၢ်ကၤကလံၤထံးထံကီၢ်တဖဉ်အပူၤလီၤ. ပုၤအါတက့ၢ်လၢ အအိဉ်ဒီး schistosomiasis တအိဉ်ဒီးတၢ်ဆါပနီဉ် နီတမံၤဒီးဘဉ်ကူဘဉ်ကတၢ်တၢ်လၢ အပူၤကွၢ်တဘျး န့ဉ်တဖဉ်န့ဉ်လီၤ. ဖဲတၢ်ဘဉ်ကူဘဉ်ကအံၤတၢ်မ့ၢ်တကူစါအီၤဘဉ်န့ဉ်ဒုးကဲ ထီဉ်သ့ဉ်, ကလံၤမ့တမ့ၢ်ဆံဉ်ဒါအတၢ်ဂ့ၢ်ကီၢ်တဖဉ်သ့ န့ဉ်လီၤ.

တချးအီကသံဉ်တခါအံၤအခါ,နကပြဲတဲဘဉ်နကသံဉ် သရဉ်မ့တမ့ၢ်ကသံဉ်သရဉ်သရဉ်မ့ဉ်ပုၤကွၢ်ဆါဖဲနမ့ၢ် အီကသံဉ်အဂၤတဖဉ်အခါတက့ၢ်. အဒိန့ဉ်,ကသံဉ်ကသီတဖဉ်လၢတၢ်ကသူအီၤလၢတၢ်က ကူစါတၢ်ထီဉ် (အဲးပလဲးစံဉ်), TB ဒီး HIV န့ဉ်ဘဉ်တံာ် တၢ်လိာ်အသးဒီးကသံဉ်ကသီတခါအံၤသ့လီၤ.

နကပြဲကတိၤစ့ၢ်ကီၢ်တၢ်ဒီးနကသံဉ်သရဉ်မ့တမ့ၢ်သရဉ် သရဉ်မ့ဉ်ကွၢ်ပုၤဆါဖဲနအိဉ်ဒီးတၢ်ထီဉ်တဖဉ်မ့တမ့ၢ်လၢ ညါမ့ၢ်အိဉ်ဒီးသး ဖျါအစံဉ်ဖုလၢတမ့ၢ်ဒ်ညီန့ၢ်အသိးတခါနခါန့ဉ်လီၤ.

မိုးယကအီ Praziquantel ဒ်လဲဉ်.

ကသံဉ်တခါအံၤအကသံဉ်ဖျါတဘျီအဂီၢ်န့ဉ်ဘဉ်တၢ် ဒွးဃာ်အီၤလၢအဒိးသန့ၤထီဉ်လၢနတယၢ်န့ဉ်လီၤ. နကလိဉ်အီကသံဉ်ခံဖျါဒီးတဖျါန့ဉ်လွံၢ်န့ဉ်ရံဉ်တဖျါ လီၤ. ကသံဉ်တခါအံၤတၢ်ကပြဲအီၤဃုာ်ဒီးတၢ်အိဉ်တၢ် အီၤလီၤ. တဘဉ်အ့ဉ်ဘျးကသံဉ်အံၤတဂ့ၢ်. တဘဉ်အီတၢ်ဆံဉ်သ့ဉ်ဖးဒိဉ်အထံဖဲနအီကသံဉ် တခါအံၤအခါတဂ့ၢ်. တဘဉ်နီဉ်သိလ့ဉ်မ့တမ့ၢ်ဒုးလဲၤတရံးစဲးဖီကဟဉ်လၢ အလီၤပျံၤဖဲအီကသံဉ်တခါအံၤအခါတဂ့ၢ်.

If you vomit soon after you take the medication, please tell your doctor as the dose may need to be repeated.

နမ့်ၵ်းဖဲအီလီၤကသံၵ်တခါအံၤဝံၤအလီၢ်ခံၵ်,ဝံၤသး
စူၤကတီၢ်တၢ်ဒီးနကသံၵ်သရၵ်မ့ၢ်လၢနကဘၣ်အိ
က့ၤကသံၵ်အံၤအယိၵ်န့ၣ်လီၤ.

Are there any side effects?

Possible side effects from this medicine may include dizziness or drowsiness.

ကသံၵ်အတၢ်ဘၣ်ဒိဘၣ်ထံးအိၣ်တမံၤ မံၤၤၤ

ကသံၵ်အတၢ်ဘၣ်ဒိလၢအအိၣ်ထီၣ်သ့န့ၣ်ဟ်ဃုာ်တၢ်
မူၤမ့တမ့ၢ်မံၣ်တဖၣ်န့ၣ်လီၤ.

Some people may also have

- headache
- nausea or vomiting
- itching
- fever.

ပုၤတနီၤဘၣ်သ့ၣ်သ့ၣ်ကအိၣ်စ့ၢ်ကီးဒီး

- ခိၣ်သ့ၣ်ဃံဆါ
- သးကလဲာ်မ့တမ့ၢ်ဘျီး
- တၢ်သး
- တၢ်ကိၢ်ထီၣ်

These symptoms should go away after a few days. Tell your doctor if these symptoms do not go away.

တၢ်ဆါပနီၣ်သ့ၣ်တဖၣ်အံၤကၢၤဂြၢ်မၢ်ဘၣ်အသးလၢ
စ့ၢ်ကိၢ်သီအတီၢ်ပူၤန့ၣ်လီၤ.

တဲဘၣ်နကသံၵ်သရၵ်ဖဲတၢ်ဆါပနီၣ်တဖၣ်အံၤမ့ၢ်တ
ဟါမၢ်ကွံာ်ဘၣ်အခါတက့ၢ်

If you develop a rash or swelling of the face, lips, tongue or other parts of the body you should tell a doctor immediately (call doctor or go straight to emergency department).

ဖဲနမ့ၢ်အိၣ်ထီၣ်ဒီးတၢ်သးမ့တမ့ၢ်တၢ်ညိးထီၣ်လၢနမဲာ်
သ့ၣ်နိးဖံး,ပူၤမ့တမ့ၢ်နနီၢ်ခိအတၢ်လီၢ်အဂၢၤန့ၣ်နကၢၤဂြၢ်
ဘၣ်နကသံၵ်သရၵ်တဘျီဃီလီၤ(ကိးဘၣ်ကသံၵ်သ
ရၵ်မ့တမ့ၢ်လဲၤလီၤလီၤဆူဂ့ၢ်ဂီၢ်အူအဝဲၤကျိၤတက့ၢ်).

Rare side effects include:

- seizures,
- heart palpitations
- chest pain.

ကသံၵ်အတၢ်ဘၣ်ဒိလၢတနီၤစိၤကဲထီၣ်တဖၣ်

- တၢ်ထီၣ်တဖၣ်,
- သးဖျါၣ်ကနိး
- သးနါပှၢ်ဆါ

If these occur, you should also tell a doctor immediately.

ဖဲတၢ်သ့ၣ်တဖၣ်အံၤမ့ၢ်ကဲထီၣ်န့ၣ်,နကၢၤဂြၢ်ဘၣ်န
ကသံၵ်သရၵ်တဘျီဃီန့ၣ်လီၤ.

More information is available for this medicine in the form of Consumer Medicines Information (CMI). CMIs are available from hospitals, pharmacies and medical practices and from the Therapeutic Goods Administration website (www.tga.gov.au/consumer-medicines-information-cmi). The CMI contains important information about your medicine and must be read in conjunction with this leaflet

တၢ်ဂ့ၢ်တၢ်ကျိၤဆူညါန့ၢ်အိၣ်ဝဲဒၣ်လၢကသံၣ်တခါ အံၤအဂီၢ်လၢက့ၢ်ဂီၤလၢအမ့ၢ် ပှၤသူတၢ်အဂီၢ်ကသံၣ်ကသီအတၢ်ဂ့ၢ်တၢ်ကျိၤ (CMI) န့ၢ်လီၤ. CMIs န့ၢ်အိၣ်ဝဲဒၣ်လၢတၢ်ဆါဟံၣ်,ကသံၣ်ကျးဒီး ကသံၣ်ဒါးလၢကသံၣ်သရၣ်မၤတၢ်အလီၢ်ဒီးလၢ တၢ်ကူစါယါဘျါအပီးအလီၢ်အတၢ်ရဲၣ်ကျဲၤပၤဆှၢ (Therapeutic Goods Administration) အပှၢယဲၤသန့ အလီၤန့ၢ်လီၤ. (www.tga.gov.au/consumer-medicines-information-cmi). CMI ဟံၣ်ယုၣ်တၢ်ဂ့ၢ်တၢ်ကျိၤလၢအကါဒိၣ်ဘၣ်ဃးန့ၢ်ကသံၣ်ကသီဒီးတၢ်ကဘၣ်ဖးအီၤလၢအအိၣ်လိၤလိၤဒီး လံာ်ဟ့ၣ်တၢ်ဂ့ၢ်တဘျီအံၤအသိးလီၤ.

The Humanitarian Entrant Health Service (HEHS)
1/311 Wellington Street Perth 6000
Phone 9222 8500
Fax: 9222 8501
Email: HEHS@health.wa.gov.au
Opening hours: 8:15-4:15
Monday to Friday
(closed weekends and public holidays)

ပှၤန့ၢ်လီၤထံၣ်ကီၢ်လၢတၢ်သူၣ်ကညီၤသးကညီၤ
အဖီခိၣ်တဖၣ်အဂီၢ် ဆူၣ်ချ့အတၢ်မၤစၢၤ (HEHS)
1/311 Wellington Street Perth 6000
လီၤတဲစိ- 9222 8500
ဖဲး(စ)- 9222 8501
အံၤမ့(လ)- HEHS@health.wa.gov.au
အိးထီၣ်အသးန့ၢ်ရံၣ်တဖၣ်- 8:15-4:15 မှၢ်ဆၣ်တုၤမ့ၢ်ဖိး
(ကးတံၢ်အသးဖဲမ့ၢ်ဘျီမ့ၢ်ဒီးဒီး
ကမ့ၢ်အနံၤသဘျီတဖၣ်န့ၢ်လီၤ.

We are proud to be a smoke-free site.



Thank you for not smoking or vaping in any buildings or on our grounds.

ပဟံၣ်ကဖၢပသးလၢပမ့ၢ်တၢ်တအိၣ်မိၣ်ထူအလီၤန့ၢ်လီၤ.



တၢ်ဘျးဒိၣ်မးလၢနတအိၣ်မိၣ်ထူ,ဆူးမိၣ်အထံလၢပတၢ်သူၣ်ထီၣ်တခါလၢလၢအပူၤအယိၤလီၤ.

This document can be made available in alternative formats on request.

© North Metropolitan Health Service 2023

လံာ်လိာ်တခါအံၤဘျီတၢ်မၤကဲထီၣ်ဃာ်အီၤလၢက့ၢ်ဂီၤအဂၤတဖၣ်အပူၤလီၤ.

© ကလံၤစိးဝှံဒိၣ်တၢ်အိၣ်ဆူၣ်အိၣ်ချ့အတၢ်မၤစၢၤ ၂၀၂၃ (North Metropolitan Health Service 2023)

